



CLINICAL UPDATE:
INITIAL REAL-WORLD
OUTCOMES OF A
COMPREHENSIVE APPROACH
TO TINNITUS MANAGEMENT
INCLUDING BIMODAL
NEUROMODULATION
(NEUROMOD LENIRE)

Craig A. Kasper, AuD




High Responder Rates With Further Improvement By Second Follow-Up




¹MCID = Minimal Clinical Important Difference (7 points decrease in the THI)
Zeman, S., Koller, M., Figueroa, R., Agresti, A., Bates, M., Coelho, C., Kleinjung, T., De Ridder, D., Langguth, B. and Landgrebe, M., 2011. Tinnitus handicap inventory for evaluating treatment effects: which changes are clinically relevant?. *Otolaryngology-Head and Neck Surgery*, 145(2), pp.282-287.

Those With More Bothersome Tinnitus Had A Higher Response Rate



¹MCID = Minimal Clinical Important Difference (7 points decrease in the THI)

Those With More Bothersome Tinnitus Had A Higher Response Rate That Further Improved With Additional Treatment



*The higher responder rate and further improvements in THI shows the benefit of patients continuing with treatment.

¹MCID = Minimal Clinical Important Difference (7 points decrease in the THI)

OVERALL HIGHLIGHTS

- FAST-ACTING RESPONSE TO TREATMENT WITH **68.1%** OF PATIENTS HAVING A MEANINGFUL IMPROVEMENT IN THI SCORES BY THE FIRST FOLLOW-UP APPOINTMENT.
- OVERALL HIGH RESPONSE RATE TO TREATMENT OF **78.3%** BY THE SECOND FOLLOW-UP APPOINTMENT.
- PATIENTS WHO WERE MODERATE OR WORSE ON THE THI SCORE AT THE ONSET OF TREATMENT HAD AN EVEN HIGHER RESPONSE RATE OF **88.9%** BY THE SECOND FOLLOW-UP VISIT.

FOR MORE INFORMATION

NYHD | INSTITUTE FOR HEARING & BALANCE
120 EAST 56TH STREET, SUITE 430
NEW YORK, NEW YORK 10022

NYHD | TINNITUS CARE
950 FRANKLIN AVENUE, LOWER LEVEL
GARDEN CITY, NEW YORK 11530

WWW.NEWYORKHEARINGDOCTORS.COM

TEL. 212 774 1971